



CLINICAL GUIDELINE

Management of Insomnia Guideline

A guideline is intended to assist healthcare professionals in the choice of disease-specific treatments.

Clinical judgement should be exercised on the applicability of any guideline, influenced by individual patient characteristics. Clinicians should be mindful of the potential for harmful polypharmacy and increased susceptibility to adverse drug reactions in patients with multiple morbidities or frailty.

If, after discussion with the patient or carer, there are good reasons for not following a guideline, it is good practice to record these and communicate them to others involved in the care of the patient.

Important Note:

The Intranet version of this document is the only version that is maintained. Any printed copies should therefore be viewed as 'Uncontrolled' and as such, may not necessarily contain the latest updates and amendments.

Original Date of Approval:	31 st July 2012		
Date of Update:	26 th April 2018		
Review Date:	30 th April 2021		
Lead Author:	L Templeton		
Approval Group:	Mental Health Quality & Clinical Governance Group		
Clinical Content Changes (Y/N):	Yes		
Version no:	2	MHS No:	35

NHS GG&C Mental Health Services Management of Insomnia Guideline

Revision Information

Please record brief details of the changes made alongside the next version number. If the procedural document has been reviewed **without change**, this information will still need to be recorded although the version number will remain the same.

Version	Date	Brief Summary of Changes	Author(s)
1.0	Apr 2015	Consultation Draft	P Davies
2.0	Apr 2018	<p>Title of guideline changed from Hypnotic Prescribing to Management of Insomnia</p> <p>Page 2,4,6. NICE CKS 2015 Insomnia added as reference</p> <p>Page 3. Sleep hygiene advice tabulated and updated</p> <p>Page 3. Choice and medication Insomnia guide and sleep hygiene handy fact sheet added</p> <p>Page 4. Potential adverse effects (including falls risk) and long term complications of hypnotics added under general principles</p> <p>Page 4,6. NICE CKS Benzodiazepine and Z-drug withdrawal 2015 added as reference</p> <p>Page 5,6. GG&C formulary site added as reference</p> <p>Page 5 Diazepam and nitrazepam not recommended due to their long half lives</p>	L Templeton L Burns

NHS GG&C Mental Health Services Management of Insomnia Guideline.

Contents

1.0	Introduction & Background	3
2.0	Scope	3
3.0	Non-pharmacological Interventions.....	3
4.0	Pharmacological Interventions	4
5.0	Choice of Hypnotic.....	5
6.0	Management of Insomnia Guideline: Audit Criteria	5
7.0	References.....	5

1.0 Introduction & Background

Insomnia can often be poorly managed therefore this guideline describes best practice around the promotion of sleep hygiene and appropriate pharmacological intervention. This guideline was originally produced by consolidating the best practice from the existing set of guidelines in use within NHS GG&C. This guideline has been updated to take into account NICE CKS Insomnia 2015¹ and NICE Technology Appraisal No77, 2004², Guidance for the short term management of insomnia.

2.0 Scope

Application – all clinical settings across Mental Health Services, NHS GG&C.

3.0 Non-pharmacological Interventions

Before treating insomnia with drugs, prescribers should consider:

- Managing any underlying cause for insomnia
- Substance misuse or dietary issues
- Ensure appropriate timing of potentially stimulating drugs
- Ensure realistic expectation of sleep and sleep duration
- Encourage sleep hygiene approaches, as indicated within the table below;

Sleep Hygiene Advice * Sleep hygiene is a variety of different practices and habits intended to promote better quality sleep
Maintain a regular sleep-wake schedule. Go to bed and rise at the same time each day
Avoid sleeping in after a poor night
Do not nap, especially close to bedtime
Keep the environment dark, quiet (use earplugs if necessary) and at a comfortable temperature
Avoid using backlit screens e.g. smart phones, tablets, computer screens, television for an hour before bedtime
Avoid watching the clock
Do not lie in bed for prolonged periods awake. If unable to fall asleep (or back to sleep) within 20 minutes, get out of bed. Engage in relaxing activity until drowsy then return to bed and repeat this action as necessary
Avoid excessive liquids or heavy evening meals
Minimise or avoid caffeine (N.B. contained in many soft drinks), alcohol, tobacco, stimulants, especially after noon
Warm milk-based drinks before bed can help. Milk contains tryptophan, a precursor to serotonin, which aids sleep.
Exercise regularly but not within 3-4 hours of bedtime
Go outside for at least half an hour each day (especially late afternoon). Exposure to natural light helps restore circadian rhythm

For further information on sleep hygiene and handy fact sheets for patients please refer to the Choice and medication website hosted via the NHS Inform portal³:

http://www.choiceandmedication.org/nhs24/condition/insomnia/Sleep_hygiene_handy_fact_sheet

If these approaches fail or are impractical, a pharmacological intervention may be appropriate.

4.0 Pharmacological Interventions

4.1 General Principles

Hypnotic use is for short term use only

- Be aware of (and advise patients of) interactions with alcohol and other drugs (prescribed and non-prescribed)
- Be aware of potential 'hangover' effects of hypnotics
- Potential adverse effects of hypnotics include daytime sedation, poor coordination, cognitive impairment, and related concerns about the increased risk of driving accidents and falls especially in the elderly
- Avoid hypnotics in patients at risk of respiratory depression
- Avoid hypnotics in patients with a severe hepatic impairment
- Avoid hypnotics in addiction-prone individuals
- As required hypnotics should not be administered routinely and ideally not at a fixed time. A clear need should be evident e.g. patient fails to sleep despite an adequate time in bed
- Long-term use of hypnotics can lead to the development of tolerance, physical or behavioural dependence, adverse effects on withdrawal, rebound insomnia, and increased mortality ¹

4.2 New presentation of a sleep disorder

- Use the recommended formulary option ⁴ i.e. zopiclone unless an alternative is clinically indicated
- Use the lowest effective dose
- Use intermittent dosing e.g. alternate nights or lower frequency of use
- Prescribe as an 'as required' medication. The prescribing and administration of regular hypnotics at a fixed time is inappropriate
- Prescription should be for short term use (no more than four weeks duration)
- Prescriptions must be reviewed regularly (at least weekly)
- If a hypnotic is commenced as an inpatient, this must be discontinued prior to patient's discharge if possible

4.3 Pre-existing prescription of a hypnotic for a sleep disorder

- Assess the current requirement for a hypnotic;
 - Evidence of an existing sleep disorder
 - Consider factors contributing to a sleep disorder
 - Appropriate dose and duration of treatment
 - Appropriate timing of dose
- If the prescription is no longer appropriate, discontinue slowly to minimise the risk of rebound insomnia and withdrawal symptoms. For further information, refer to the NICE CKS on benzodiazepine and z-drug withdrawal⁵ or seek advice from a specialist mental health pharmacist
- If a hypnotic is still required
 - Ensure formulary choice
 - Switch if necessary and optimise dose
- Prescriptions must be reviewed regularly (at least weekly)
- At resolution of sleep disorder, consider slow withdrawal of treatment

5.0 Choice of Hypnotic

From NICE Technology Appraisal No77 2004²;

“It is recommended that, because of the lack of compelling evidence to distinguish between zaleplon, zolpidem, zopiclone or the shorter-acting benzodiazepine hypnotics, the drug with the lowest purchase cost (taking into account daily required dose and product price per dose) should be prescribed.”

Patients who have not responded to one of these hypnotic drugs should not routinely be prescribed any of the others. Although, consideration may be given to alternatives in situations where there is an incidence of z-drug or benzodiazepine abuse, where alternatives may be appropriate.

The Formulary Preferred List hypnotic in NHSGG&C is zopiclone.⁴

If zopiclone is inappropriate, another formulary hypnotic should be prescribed.^{4,6} Diazepam and nitrazepam are not recommended because their long half-life commonly gives rise to next day residual effects, and repeated doses tend to be cumulative.¹

6.0 Management of Insomnia Guideline: Audit Criteria

Criterion Statement	Standard	Exceptions
Is there a documented assessment indicating a sleep disorder	100%	None
Non-pharmacological interventions have been attempted	100%	None
Where a hypnotic is indicated, zopiclone is prescribed	100%	Intolerance or previous/potential misuse of zopiclone Prescriber has documented rationale for alternative choice of hypnotic.
Evidence of regular (weekly) review	100%	None
New hypnotic prescriptions are discontinued prior to discharge if commenced as an inpatient	100%	None

7.0 References

1. National Institute for Health and Care Excellence (NICE) Clinical Knowledge Summaries (CKS). Insomnia. April 2015 <https://cks.nice.org.uk/insomnia> Accessed online 21st February 2018
2. National Institute for Health and Care Excellence (NICE) Technology Appraisal No 77. Guidance on the use of zaleplon, zolpidem and zopiclone for the short-term management of insomnia. April 2004 (reviewed 2010) <https://www.nice.org.uk/guidance/ta77> Accessed online 21st February 2018.
3. NHS Choice and Medication <http://www.choiceandmedication.org/nhs24/>
4. NHS GG&C Formulary. <http://www.ggcmedicines.org.uk/>
5. National Institute for Health and Care Excellence (NICE) Clinical Knowledge Summaries (CKS). Benzodiazepine and Z-drug withdrawal. April 2015 <https://cks.nice.org.uk/benzodiazepine-and-z-drug-withdrawal> Accessed online 21st February 2018
6. British National Formulary (BNF) eBNF